

# Pig Performance Independent of Wheat Class

Ruurd T. Zijlstra, David Overend<sup>1</sup>, David R. Hickling<sup>2</sup>, P. Howard Simmins<sup>3</sup>, and John F. Patience



## SUMMARY

Feed processing and nutritional quality for CPS and Durum wheat have traditionally been expected to be lower than for Hard Red Spring (HRS). Performance of weaned pigs was compared among six wheat classes, whilst considering particle size and diet pellet quality. Results indicated that feed processing quality and growth performance did not differ among wheat classes. Weaned pigs fed various classes of wheat including CPS and Durum may grow similarly.



## INTRODUCTION

The processing and nutritional quality of wheat is expected to vary among classes; CPS and Durum wheat are currently segregated. A range in wheat protein and fibre or non-starch polysaccharide (NSP) content may partly cause quality variation. The present study was designed to test whether wheat class by itself impacts feed processing or nutritional quality.

## EXPERIMENTAL PROCEDURES

Two cultivars from each of six classes (CPS White and Red, HRS, Durum, Hard Red Winter (HRW) and Hard White (HW)) were collected (Table 1). Protein ranged from 12.2 to 17.4% and total NSP from 9.0 to 11.5% (Table 2). A 3-week growth and digestibility study was conducted with 12-kg weaned pigs (PIC; 39-day-old; 4 pigs/pen, 12 pens per cultivar), which were fed pelleted 65%-wheat diets (3.5 Mcal DE/kg; 3.4 g dig. lysine/Mcal).

*“CPS and Durum wheat can be processed and fed to weaned pigs without compromising growth performance”*

## RESULTS

Wheat particle size ranged from 536 to 734  $\mu\text{m}$  (10/64"-screen) (Table 2). Pellet Durability Index was 96 for all diets. Feed processing quality was thus excellent for all wheat classes. In the growth study, average daily gain (ADG), average daily feed intake (ADFI), and feed efficiency did not differ among wheat classes for day 0 to 21 (Figure 1). However, some minor differences were observed in the first week. For example, ADG for Durum was 9% lower than for HRW, and similar among other classes; ADFI for HW was 7% lower than for HRW, and similar among other classes. Finally, diet energy digestibility (and thus DE content) was lowest for CPS Red (86.5%), medium for CPS White, HRS and HW (87.2 to 87.5%) and highest for HRW and Durum (88.6 & 88.9%) (Figure 2).

<sup>1</sup> Ridley Inc., Mankato, MN

<sup>2</sup> Canadian International Grains Institute, Winnipeg, MB; presently Canola Council of Canada, Winnipeg, MB

<sup>3</sup> Danisco Animal Nutrition, Marlborough, UK

## IMPLICATIONS

Protein but not NSP content varied among 12 wheat cultivars harvested in western Canada in 2001. Wheat protein content was “corrected for” during diet formulation and did not affect pig performance. Wheat NSP content was low overall, indicating that all 12 wheat samples were of excellent nutritional quality. Still, DE content ranged 7% and was highest for Durum. Reductions in ADG and ADFI for CPS and Durum wheat were limited to the first two weeks, and did not exist after 3 weeks. In conclusion, despite variations in wheat DE content, weaned pigs fed various classes of wheat including CPS and Durum performed effectively, and wheat class by itself was not a cause for a range in feed processing or nutritional quality.

Canadian Wheat Board Class	Popular Name
Canadian Prairie Spring White	CPS White
Canadian Prairie Spring Red	CPS Red
Canadian Western Red Spring	Hard Red Spring (HRS)
Canadian Western Amber Durum	Durum
Canadian Western Red Winter	Hard Red Winter (HRW)
Canadian Western Hard White	Hard White (HW)

Wheat class	Protein (% as is)	Total NSP (% as is)	Particle size ( $\mu$ m)
CPS White	15.1, 16.1	11.3, 11.4	591, 700
CPS Red	12.4, 15.8	11.0, 11.3	556, 631
HRS	15.4, 17.1	10.8, 11.5	640, 708
Durum	16.3, 16.8	9.0, 10.1	624, 734
HRW	12.2, 13.7	9.7, 10.9	536, 636
HW	16.5, 17.4	10.9, 11.3	629, 724

## FINANCIAL IMPLICATIONS

Typically feed quality durum wheat can be purchased at a discount relative to other wheat types. Therefore producers have one more opportunity to reduce their feed cost per pig without compromising growth performance.

## ACKNOWLEDGEMENTS

Strategic program funding provided by Sask Pork, Alberta Pork, Manitoba Pork, and Saskatchewan Agriculture and Food Development Fund. Project funding was provided by the Canadian Wheat Board, Canadian International Grains Institute, Feed-Rite (Ridley Inc.), Danisco Animal Nutrition, Degussa, and Quality Assured Seeds.




Figure 1 Effects of wheat class on average daily gain (ADG) and feed intake (ADFI) of weaned pigs in the 3-week growth study.




Figure 2 Effect of wheat class on diet energy digestibility measured in weaned pigs.



**Prairie Swine Centre**  
 Box 21057  
 2105 8<sup>th</sup> Street East  
 Saskatoon, Saskatchewan  
 S7H 5N9  
 Phone (306) 373-9922  
 Fax (306) 955-2510  
 E-mail whittington@sask.usask.ca

Prairie Swine Centre is an affiliate of



<sup>1</sup>Ridley Inc., Mankato, MN

<sup>2</sup>Canadian International Grains Institute, Winnipeg, MB; presently Canola Council of Canada, Winnipeg, MB

<sup>3</sup>Danisco Animal Nutrition, Marlborough, UK